



HYPERBOLIC SPACE GROUPS FOUR SERIES WITH SIMPLICIAL DOMAINS

Dr. Avinash Kumar Yadav

Email: avinashyad75@gmail.com

Associate Professor, Department Of Applied Science

Greater Noida College Of Technology, Greater Noida, India

Abstract:

Hyperbolic space groups are isometric groups, acting discontinuously on the hyperbolic 3-space with compact fundamental domain. One possibility to classify them is to look for fundamental domains of these groups. Here are considered super groups for four series of groups with simplified fundamental domains. Considered simplices, denoted in [9] by T_{19} , T_{46} , T_{59} , belong to family F12, while T_{31} belongs to F27.

1. Introduction

Hyperbolic space groups are isometric groups, acting discontinuously on the hyperbolic 3-space with compact fundamental domain. One possibility to classify them is to look for fundamental domains of these groups. Face pairing identifications of a given polyhedron give us generators and relations for a space group by Poincare Theorem [1], [3], [7].

The simplest fundamental domains are simplices and truncated simplices by polar planes of vertices when they lie out of the absolute. There are 64 combinatorial different face pairings of fundamental simplices [16], [6], furthermore 35 solid transitive non-fundamental simplex identifications [6]. I. K. Zhuk [16] has classified Euclidean and hyperbolic fundamental simplices of finite volume up to congruence. Some completing cases are discussed in [2], [5], [10], [12], [13], [14], [15]. Algorithmic procedure is given by E. Molnar and I. Prok [5]. In [6], [8] and [9] the authors summarize all these results, arranging identified simplices into 32 families. Each of them is characterized by the so-called maximal series of simplex tiling's. Besides spherical, Euclidean, hyperbolic realizations there exist also other metric realizations in 3-dimensional simply connected homogeneous Riemannian spaces, moreover, metrically non-realizable topological simplex tiling's occur as well [4].

When vertices are out of the absolute, the simplex is not compact and then we truncate it with polar planes of the vertices. The new compact polyhedron obtained in that way, let us call it trunc-simplex, is fundamental domain of some larger group. It has new triangular faces whose pairing gives new generators. For simplicity, here we require that the new pairing generators keep the original simplicial face structure. Other possibilities will be discussed elsewhere. Dihedral angles around new edges are $\frac{1}{4}\pi=2$. That means that there will be four congruent polyhedra around them in a new fundamental space tiling. These investigations have been initiated by the author (see e.g. [14]).

Each identified simplex, considered in this paper, has two equivalence classes for edges with three edges in each. Edges in the same class haven't common vertex. There are 4 different face pairings: T_{19} , T_{46} , T_{59} in family F12 and T_{31} in family F27 to investigate in this paper to extend the series tabled in [9].

In Section 2 we recall Poincare Theorem which provides a method to construct discontinuously acting isometric groups.

2. Construction of discontinuously acting isometric groups

In other words the segment e_1 is successively surrounded by polyhedral

$P; P g_1^{i1}; P g_2^{i1} g_1^{i1}; \dots; P g_r^{i1} g_2^{i1} g_1^{i1}$ which will an angular region of measure $2\pi/4 = \pi/2$. In the above case 1. the following holds

(2.2) $"(e_1) + \text{ccc} + "(e_i) + "(e_{i+1}) + \text{ccc} + "(e_{i+1+j}) = \pi/4 = \pi/2$: In case 2. we have

(2.3) $"(e_1) + \text{ccc} + "(e_r) = \pi/4 = \pi/2$: Finally, the cycle transformation $c = g_1 g_2 : : : g_r$ belonging to the edge segment class $fe_1 g$ is a rotation, say, of order n . Thus we have the cycle relation in both cases (2.4) $(g_1 g_2 : : : g_r)^n = 1$:

Throughout in this paper we shall apply the specified Poincare theorem:

Theorem 2.1. Let P be a polyhedron in a space S^3 of constant curvature and G be the group generated by the face identifications, satisfying conditions (a)-(c). Then G is a discontinuously acting group on S^3 , P is a fundamental domain for G and the cycle relations of type (2.4) for every equivalence class of edge segments form a complete set of relations for G , if we also add the relations $g_i^2 = 1$ to the occasional involutive generators $g_i = g_i^{i1}$.

3. Isometry groups of simplices and their super groups

3.1. SIMPLEX T_{19}

Face pairing isometries for simplex T_{19} (6a; 6b) (Figure 1) are

$$r_0: \begin{matrix} A_1 A_2 A_3 \\ A_3 A_2 A_1 \end{matrix} ; r_1: \begin{matrix} A_0 A_2 A_3 \\ A_2 A_0 A_3 \end{matrix} ; r_2: \begin{matrix} A_0 A_1 A_3 \\ A_3 A_1 A_0 \end{matrix} ; r_3: \begin{matrix} A_0 A_1 A_2 \\ A_0 A_2 A_1 \end{matrix} ;$$

Relations for the isometry group are obtained by Theorem 2.1 and the presentation

$$i(T_{19}; 6a; 6b) = \langle r_0, r_1, r_2, r_3 \mid r_0^2 = r_1^2 = r_2^2 = r_3^2 = 1, (r_0 r_1 r_2 r_1 r_0 r_3)^a = (r_3 r_2 r_0 r_2 r_3 r_1)^b = 1; a, b \in \mathbb{N} \rangle;$$

Considering vertex figures on a symbolic 2-dimensional surface (plane) around the vertices, we can glue a fundamental domain for the stabilizer subgroup, e.g. $i(A_2)$ of vertex A_2 . Transformation r_1 maps vertex A_2 onto A_0 and T_{A_2} onto $T_{A_0}^{r_1}$. That means that T_{A_2} and $T_{A_0}^{r_1}$ have a joint edge corresponding to the joint face f_{r_1} of simplex T . Similarly, vertex figures T_{A_2} and $T_{A_1}^{r_3}$ have joint edge corresponding to f_{r_3} , and $T_{A_1}^{r_3}$ and $T_{A_3}^{r_0 r_3}$ to $(f_{r_0})^{r_3}$. One fundamental domain for $i(A_2)$ (Figure 2) is $P_{A_2} := T_{A_0}^{r_1} \cup T_{A_2} \cup T_{A_1}^{r_3} \cup T_{A_3}^{r_0 r_3}$

Figure 1. The simplex T_{19}




Figure 2. The fundamental domain P_{A_2} for i_{A_2} and the generators for i_{A_2} , obtained from P_{A_2} , are $r_3 r_0 r_2 r_1 : (f_{r_2})^{r_0 r_3} ! (f_{r_2})^{r_1}$; $r_0 : f_{r_0} ! f_{r_0}$; $r_1 r_3 r_1 : (f_{r_3})^{r_1} ! (f_{r_3})^{r_1}$; $r_3 r_2 r_3 : (f_{r_2})^{r_3} ! (f_{r_2})^{r_3}$; $(r_3 r_0) r_1 (r_0 r_3) : (f_{r_1})^{r_0 r_3} ! (f_{r_1})^{r_0 r_3}$:

In the diagram for P_{A_2} the minus sign in notations a^i , b^i means that edges in these classes are directed to the considered vertex, (the plus sign in diagram means the opposite direction).

When parameters a , b are large enough, namely $1=a+1=b<2$, by angle sum criterion for P_{A_2} , then simplex T is hyperbolic with the vertices out of the absolute [9]. Then it is possible to truncate the simplex by polar planes of these vertices. In such a way we get a compact trunc simplex (with 8 faces) denoted by $O_{19}(6a; 6b)$. If we equip O_{19} with additional face pairing isometries, it will be a fundamental domain for a group $i_j(O_{19}; 6a; 6b)$ which will be a super group of $i(T_{19}; 6a; 6b)$. We require, also later on, that the new generators keep the original simplex face structure. A trivial group extension with plane rejections $\bar{m}_i$, $i = 0; 1; 2; 3$, in polar planes of the outer vertices A_i is always possible (Figure 3). Then the new group, by Theorem 2.1 is $i_1(O_{19}; 6a; 6b) = (r_0; r_1; r_2; r_3; \bar{m}_0; \bar{m}_1; \bar{m}_2; \bar{m}_3 ; i_1 r_0^2 = r_1^2 = r_2^2 = r_3^2 = \bar{m}_0^2 = \bar{m}_1^2 = \bar{m}_2^2 = \bar{m}_3^2 = (r_0 r_1 r_2 r_1 r_0 r_3)^a = (r_3 r_2 r_0 r_2 r_3 r_1)^b = \bar{m}_0 r_3 \bar{m}_0 r_3 = \bar{m}_1 r_2 \bar{m}_1 r_2 = \bar{m}_2 r_0 \bar{m}_2 r_0 = \bar{m}_3 r_1 \bar{m}_3 r_1 = \bar{m}_0 r_2 \bar{m}_3 r_2 = \bar{m}_1 r_3 \bar{m}_2 r_3 = \bar{m}_0 r_1 \bar{m}_2 r_1 = \bar{m}_1 r_0 \bar{m}_3 r_0 = 1)$:




Figure 3. The trunc simplex O_{19}^1 with trivial group extension

There is a further possibility to equip the new triangular faces with face pairing isometries (Figure 4). New additional face pairings of O_{19} have to satisfy the follow-ing criteria. Polar plane of A_2 and so stabilizer i_{A_2} will be invariant under these new transformations, Xing A_2 , and exchanging the half spaces obtained by the polar plane. Thus, fundamental domain P_{A_2} is divided into two parts, and the new stabilizer of the polar plane will be a super group for i_{A_2} , namely of index two. Inner symmetries of the P_{A_2} -tiling give us the idea how to introduce a new generators. Let g be the glide reflection as a composition of the translation in the plane of the vertex Figure with a rejection in this plane. Then g maps the vertex Figure T_{A_2} onto $T_{A_3}^{r_0 r_3}$

and $T_{r_0 r_3}^{A_3}$ onto $T_{r_1 r_2 r_0 r_3}^{A_2}$, equivalent to $T_{A_2}^{A_0}$. Then g also maps $T_{r_1}^{A_0}$ onto $T_{r_3}^{A_1}$ and $T_{r_3}^{A_1}$ onto $T_{r_2 r_0 r_3}^{A_0}$,

equivalent to $T_{r_1}^{A_0}$. In that case the new generators for $i_2(O_{19}; 6a; 6b)$ will be g and $g = r_1 g r_1$ in Figure 4,

while the new group, by Theorem 2.1 is $i_2(O_{19}; 6a; 6b) = (r_0; r_1; r_2; r_3; g_1; g_2 ; i_2 r_0^2 = r_1^2 = r_2^2 = r_3^2 = (r_0 r_1 r_2 r_1 r_0 r_3)^a = (r_3 r_2 r_0 r_2 r_3 r_1)^b = r_3 g_1 r_2 g_1^{i_1} = g_1 r_3 g_2 r_2 = g_1 r_0 g_2^{i_1} r_1 = r_0 g_2 r_1 g_2^{i_1} = 1)$:

The P_{A_2} -tiling in the polar plane of A_2 do not allow other identifications on the truncated simplex O_{19} .




Figure 4. The trunc simplex O_{19}^2 with non-trivial group extension

Fundamental domains T_{19} and O_{19}^j ($j = 1; 2$) above, allow to divide them to smaller polyhedra, equipped with face pairing identifications. Namely, there is a half-turn $h \in A_0 A_1 A_2 A_3 A_1 A_0 A_3 A_2$ leaving invariant the tessellations of space with T_{19} or O_{19}^j , so groups $\langle T_{19}; 6a; 6b \rangle$ and $\langle O_{19}; 6a; 6b \rangle$ are not maximal. The automorphism groups ${}^2_{2i6}(3u; 3v)$ of their tilings ([8], [9]) have domains which are fundamental polyhedra of piecewise linear bent faces. That domains are obtained by identifying equivalent points, under symmetry h , of simplex T_{19} (Figure 5), and consequently also each trunc simplex O_{19}^j above ($j = 1; 2$). Since $r_3 = hr_2h$ and $r_1 = hr_0h$, presented for $a = b$, maximal groups are now (with $u = 2a$ and $v = 2b$ for the rotational parameters) by ${}^2_{2i6}(3u; 3v) = \langle h; r_0; r_2; h^2 = r_0^2 = r_2^2 = (r_0hr_0hr_2h)^u = (r_2hr_2r_0)^v = 1; u = 2a; v = 2b \rangle$ and $\langle Q; 3u; 3v \rangle = \langle h; r_0; r_2; m_1; m_2; h^2 = r_0^2 = r_2^2 = m_1^2 = m_2^2 = (r_0hr_0hr_2h)^u = (r_2hr_2r_0)^v = m_1r_2m_1r_2 = m_2r_0m_2r_0 = m_1r_2m_2r_2 = m_1r_0m_2r_0 = 1; u = 2a; v = 2b \rangle$:




Figure 5. The fundamental domain of supergroup ${}^2_{2i6}(3u; 3v)$

If $a = b$ then simplex T and trunc simplex O^j have more symmetries. Then the maximal supergroup for $\langle T_{19}; 6a; 6b \rangle$ is a Coxeter group, by [9], while the maximal supergroup for $\langle O_{19}; 6a; 6b \rangle$ might have only the trivial extension, so it is also a Coxeter group.

3.2. SIMPLEX T_{46}

For $T_{46}(6a; 3b)$, the face pairing isometries are (Figure 6):

$$\begin{array}{ccccccc} A_0 & A_1 & A_3 & & A_0 & A_1 & A_2 \\ r_2 : \bar{A} A_3 & A_1 & A_0 & !; & r_3 : \bar{A} A_0 & A_2 & A_1 !; s : \bar{A} A_2 & A_3 & A_0 !; \end{array}$$

and the tiling group is

$$\langle T_{46}; 6a; 3b \rangle = \langle r_2; r_3; s; r_2^2 = r_3^2 = (s^2 r_2 s^2 r_3)^a = (r_2 s r_3)^b = 1; a, b \in \mathbb{N} \rangle$$

One fundamental domain for the stabilizer group $\langle A_2 \rangle$ of the vertex A_2 (Figure 6) is $P_{A_2} := T_{A_0}^{r_2 s^2} [T_{A_3}^{s^2} [T_{A_2} [T_{A_1}^{r_3} [T_{A_0}^{r_2}]]]]$ and the generators are then $sr_2 r_3 r_2 s^2 : (fr_3)^{r_2 s^2} ! (fr_3)^{r_2 s^2} ; s^2 r_2 s^2 : (fs^2)^{r_2 s^2} ! (fs)^{r_2 s^2} ; r_3 s : (fs_1)^{r_3} ! fs; r_3 r_2 r_3 : (fr_2)^{r_3} ! (fr_2)^{r_3} : The stabilizer \langle A_2 \rangle of P_{A_2} above is hyperbolic i.e. (again by the angle sum criterion for P_{A_2}) 2\pi + 1 = a < 2. Then truncating the simplex by polar planes of the vertices,$




Figure 6. The simplex T_{46} and the fundamental domain P_{A_2}

a new trunc-simplex O_{46} may have plane reflections as face pairing isometries of the new faces. In this case the new group is (Figure 7)

$$\langle O_{46}; 6a; 3b \rangle = \langle r_2; r_3; s; m_0, m_1, m_2, m_3; r_2^2 = r_3^2 = m_0^2 = m_1^2 = m_2^2 = m_3^2 = (s^2 r_2 s^2 r_3)^a = (r_2 s r_3)^b = m_0 r_3 m_0 r_3 = m_1 r_2 m_1 r_2 = m_2 r_3 m_2 r_3 = m_3 r_2 m_3 r_2 = m_2 s m_3 s^2 = m_3 s m_0 s^2 = m_1 s m_2 s^2 = 1 \rangle$$




Figure 7. The trunc-simplex O_{46}

Other possibility, by symmetries of the fundamental domain P_{A_2} is the group extended by the point rejection z , indicated in Figure 6. This point rejection z (say) maps the triangle of A_2 to that of $A_3^{s_1}$ and triangle of $A_1^{r_3}$ to that of $A_0^{r_2 s_1}$ in P_{A_2} (Figure 6). Thus, the above z induces new generators g_1 and g_2 as glide reflections, pairing the truncations at A_2, A_3 and those at A_1, A_0 , respectively.

$$i_2(O_{46}; 6a; 3b) = (r_2; r_3; s; g_1; g_2; i; r_2^2 = r_3^2 = (s^2 r_2 s^2 r_3)^a = (r_2 s r_3)^b = r_2 g_2 r_3 g_2^{i^1} = g_2 r_2 g_1^{i^1} r_3 = s g_1 s g_2^{i^1} = g_1 s^{i^1} g_1 s^{i^1} = 1):$$

If r_0 and h are similarly introduced, as in the previous section, so that $r_3 = h r_2 h$ and $s = r_0 h$ hold. Then the maximal group ${}^2_{2i6}(3u; 3v)$, now with $u = 2a, v = b$, will be supergroup of $i(T_{46}; 6a; 3b)$, and $i(Q; 3u; 3v)$ extends $i_j(O_{46}; 6a; 3b)$ ($j = 1; 2$) as well.

3.3. SIMPLEX T_{59}

In the case of the simplex $T_{59}(3a; 3b)$ the face pairing identifications are (Figure 8)

$$\begin{array}{ccc} A_1 & A_2 & A_3 \\ s_1 : \bar{A}_2 & A_3 & A_0 \end{array} ; \quad \begin{array}{ccc} A_0 & A_1 & A_3 \\ s_2 : \bar{A}_2 & A_0 & A_1 \end{array} !$$

and the presentation of the group is

$$i(T_{59}; 3a; 3b) = (s_1; s_2; i; s_1^2 = s_2^2 = (s_1^2 s_2^2)^a = (s_2^2 s_1^2)^b = 1; a; b \in \mathbb{N}):$$

The stabilizer group $i(A_0)$ has fundamental domain (Figure 8)

$$\begin{array}{ccc} s & & s_i^1 \\ 2 & s_2 & 2 \end{array} \quad \text{PA0} := \text{TA } 3 \mid \text{TA } 1 \mid \text{TA0} \mid \text{TA } 2$$

and the generators

$$\begin{array}{ccc} 2 & s^2 & 1 \quad i \quad 1 \quad s \quad 1 \quad 2 \quad s_i^1 \quad s^2 \\ (f_{s_1}) & & : (f_{s_i^1}) \\ s_2 \quad s_1 & : (f_{s_i^1})^2 \quad f_{s_1}; & s_2 \quad s_1^2 s_2 : (f_{s_i^1})^2 \quad 2 \quad 2 \quad ; \quad s_2 s_1 s_2 \quad 2 \quad (f_{s_1})^2 : \\ 1 & ! & 1 \quad ! \quad 1 \quad ! \end{array}$$




Figure 8. The simplex T_{59} and the fundamental domain P_{A_0}

There are two possibilities for the isometry group with trunc-simplex O_{59} as a fundamental domain, $i \otimes 1 = a + 1 = b < 1$.

In the trivial case, group is (Figure 9)




Figure 9. The trunc-simplex O_{59}

$i_1(O_{59}; 3a; 3b) = (s_1; s_2; m_0; m_1; m_2; m_3; i; m_0^2 = m_1^2 = m_2^2 = m_3^2 = (s_1^2 s_2)^a = (s_2^2 s_1^{-1})^b = m_2 s_1 m_3 s_1^{-1} = m_3 s_1 m_0 s_1^{-1} = m_1 s_1 m_2 s_1^{-1} = m_3 s_2 m_1 s_2^{-1} = m_1 s_2 m_0 s_2^{-1} = m_0 s_2 m_2 s_2^{-1} = 1)$:

Taking g_1 and $g_2 = s_1^{-1} g_1 s_1^{-1}$ as a new generators, other possibility for the group is $i_2(O_{59}; 3a; 3b) = (s_1; s_2; g_1; g_2; i; (s_1^2 s_2)^a = (s_2^2 s_1^{-1})^b = g_1 s_1 g_2 s_1 = g_1 s_2 g_1 s_2 = s_2 g_2 s_2 g_1^{-1} = g_2 s_1^{-1} g_2 s_1^{-1} = 1)$:

Since, it is possible to express the face pairing isometries s_1 and s_2 of T_{59} by h, r_0, r_2 : $s_1 = r_0 h$ and $s_2 = r_2 h$, the groups ${}^2_{2i6}(3u; 3v)$ and $i(Q; 3u; 3v)$ are super groups of the groups $i(T_{59}; 3a; 3b)$ and $i_j(O_{59}; 3a; 3b)$, ($u = a; v = b$).

3.4. SIMPLEX T_{31}

The face pairings identifications for the simplex $T_{31}(6a; 12b)$ are (Figure 10)

$$\begin{array}{ccc} A_1 & A_2 & A_3 \\ m : \bar{A}_1 & A_2 & A_3 \\ !; r : \bar{A}_2 & A_0 & A_3 \\ !; s : \bar{A}_1 & A_3 & A_0 \end{array}$$

The group presentation is $i(T_{31}; 6a; 12b) = (m; r; s; i; r^2 = m^2 = (rmrs^{-1}ms)^a = (rs^2ms^{-1}rs^2ms^{-1})^b = 1; a, 1; b, 1)$: For the stabilizer group $i(A_1)$ one of the fundamental domains is (Figure 10) $P_{A_1} := T_A^{s_2^2} [T_A^{s_0} [T_{A_1} [T_A^{s_1^{-1}}]$




Figure 10. The simplex T_{31} and the fundamental domain P_{A_1} with generators

$$\begin{array}{ccc} srs^{-1} : (f)^{s_1^{-1}} & ! (f)^{s_1^{-1}} & ; \\ r & r & \\ s^{-1}rs : (f)^{s_2} & ! (f)^s & \\ r & r & \end{array}$$

After truncating the simplex by the polar planes of the vertices, $i \otimes 1 = b + 1 = a < 4$ trunc simplex O_{31} may have only trivial group extension (Figure 11)

$i(O_{31}; 6a; 12b) = (m; r; s; r^2 = m^2 = (rmrs^{i1}ms)^a = (rs^2ms^{i2}rs^2ms^{i2})^b = m_3rm_3r = m_0rm_2r = m_1mm_1m = m_2mm_2m = m_3mm_3m = m_1sm_3s^{i1} = m_2sm_0s^{i1} = m_0sm_1s^{i1} = 1; a, 1; b, 1)$:




Figure 11. The trunc-simplex O_{31}

It is not possible to extend generators of T_{31} by h , since then a new rejection plane on halfturn axis r would yield $a = b$ and we got the richer family F.1.



References

- [1] B. Maskit, On Poincaré's theorem for fundamental polygons, *Adv. in Math.* 7 (1971), 219{230.
- [2] E. Molnár, Eine Klasse von hyperbolischen Raumgruppen, *Beiträge zur Algebra und Geometrie* 30 (1990), 79{100.
- [3] E. Molnár, Polyhedron complex with simply transitive group actions and their realizations, *Acta Math. Hung.* 59, (1-2) (1992), 175{216.
- [4] E. Molnár, The projective interpretations of the eight 3-dimensional homogeneous geometries, *Beiträge zur Algebra und Geometrie (Contributions Alg. Geom.)* 38, Vol.2 (1997), 261{288.
- [5] E. Molnár and I. Prok, A polyhedron algorithm for finding space groups, *Proceedings of Third Int. Conf. On Engineering Graphics and Descriptive Geometry*, Vienna 1988, No.2, 37{44.
- [6] E. Molnár and I. Prok, Classification of solid transitive simplex tilings in simply connected 3-spaces, Part I, *Colloquia Math. Soc. János Bolyai* 63. *Intuitive Geometry*, Szeged (Hungary), 1991. North-Holland (1994), 311{362.
- [7] E. Molnár, I. Prok, Hyperbolic space form with 'Schläöi solid' (8; 8; 3), *Symmetry: Culture and Science, Tessellation* 35, (1-2) (2011), 263{267.
- [8] E. Molnár, I. Prok and J. Szirmai, Classification of solid transitive simplex tilings in simply connected 3-spaces, Part II, *Periodica Math. Hung.* 35, (1-2) (1997), 47{94.
- [9] E. Molnár, I. Prok and J. Szirmai, Classification of tile-transitive 3-simplex tilings and their realizations in homogeneous spaces, *Non-Euclidean Geometries, János Bolyai Memorial Volume*, Editors: A. Prékopa and E. Molnár, *Mathematics and Its Applications*, Vol. 581, Springer (2006), 321{363.
- [10] M. Stojanović, Some series of hyperbolic space groups, *Annales Univ. Sci. Budapest, Sect. Math.* 36 (1993), 85{102.
- [11] M. Stojanović, Hyperbolic realizations of tilings by Zhuk simplices, *Matematički Vesnik*, 49 (1997), 59{68.
- [12] M. Stojanović, Hyperbolic space groups for two families of fundamental simplices, *Novi Sad J. Math.*, 29, No.3 (1999), 337{348, XII Yugoslav Geometric Seminar, Novi Sad, Octobar 8-11, 1998.
- [13] M. Stojanović, Fundamental simplices with outer vertices for hyperbolic groups, *Filomat* 24/1 (2010), 1{19.
- [14] M. Stojanović, Coxeter groups as automorphism groups of solid transitive 3-simplex tilings, man-uscript.
- [15] I. K. Zhuk, Fundamental tetrahedra in Euclidean and Lobachevsky spaces, *Soviet Math. Dokl.* 270, No.3 (1983), 540{543}.