



A DOUBLE – SEQUENCE HYBRID S -ITERATION SCHEME FOR FIXED POINTS OF CONTINUOUS PSEUDOCONTRACTIONS

S. Gopinath¹, J. Gnanaraj² and S. Lalithambigai³

¹Department of Mathematics, Kamaraj College of Engineering and Technology,
Virudhunagar, Tamilnadu, India

²Department of Mathematics, Goverment Arts College, Paramakudi, Tamilnadu, India

³School of Mathematics, Madurai Kamaraj University, Madurai, Tamilnadu, India

ABSTRACT

Let C be a bounded closed convex nonempty subset of a real Hilbert space H , It is proved that a double – sequence Hybrid S- Iteration scheme converges to a fixed point of continuous pseudocontractive map T which map C into C .

Keywords : Continuous pseudocontraction, Double – sequence Hybrid S- Iteration, Hilbert space, Nonexpansive, Strong convergence.

2010 Mathematics Subject Classification: 47H09, 47H10, 47H17.

I. INTRODUCTION

Let H be a real Hilbert space and let C be a nonempty convex subset of a real Hilbert space H .

Let T be a mapping.

The following Definitions have been studied widely and deeply by many authors; see, e.g., [1-21] for more details.

Definition 1.1. The mapping T is said to be nonexpansive if

$$\|Tx - Ty\| \leq \|x - y\|, \quad \forall x, y \in C.$$

Definition 1.2. The mapping T is said to be pseudocontractive if

$$\langle Tx - Ty, x - y \rangle \leq \|x - y\|^2, \quad \forall x, y \in C.$$

Definition 1.3. Let $\mathbb{N}$ denote the set of all the nonnegative integers and let E be a normed linear space. By a double sequence in E is meant by function $f : \mathbb{N} \times \mathbb{N} \rightarrow E$ defined by $f(n, m) = x_{n,m} \in E$. The double sequence $\{x_{n,m}\}$ is



said to converges stongly to x^* if any $\varepsilon > 0$ there exist integer $N, M > 0$ such that $\forall n \geq N, m \geq M$, we have that $\|x_{n,m} - x^*\| < \varepsilon$. If $\forall n, r \geq N$ and $m, t \geq M$, we have that $\|x_{n,r} - x_{m,t}\| < \varepsilon$, then the double sequence is said to be Cauchy. Furthermore, if for each fixed n , $x_{n,m} \rightarrow x_n^*$ as $m \rightarrow \infty$ and then $x_n^* \rightarrow x^*$ as $n \rightarrow \infty$, then $x_{n,m} \rightarrow x^*$ as $n, m \rightarrow \infty$.

In the last few years or so, Several iteration process have been established for the constructive approximation of solution to several classes of (nonlinear) operator equations and several convergence results established using these iterative processes (see, e.g [1-21] and the reference cited therein). Most of these convergence results for the iterative solution of nonlinear operator equations or approximation of fixed points of nonlinear maps have used iteration process of the Mann, Ishikawa and Hybrid S- types.

The concept of Mann-type double –sequence iteration process introduced by C. Moore and he prove that it converges strongly to a fixed point of a continuous pseudocontraction which maps a bounded closed convex nonempty subset of a real Hilbert space into itself.

In this paper, we prove the same result for double –sequence of Hybrid S-iterative scheme.

II. MAIN RESULTS

Theorem 2.1. Let C be a bounded closed convex nonempty subset of a real Hilbert space H , and $T: C \rightarrow C$ be a continuous pseudocontractive map. Let $S: C \rightarrow C$ be nonexpansive mapping satisfying that $\|x - Sy\| \leq \|Sx - Sy\|$ for all $x, y \in C$ and $\{\beta_n\}_{n \geq 0}, \{\alpha_k\}_{k \geq 0} \subset (0, 1)$ be real sequence satisfying the following condition :

- 1) $\lim_{k \rightarrow \infty} \alpha_k = 1$ (monotonically);
- 2) $\lim_{k, j \rightarrow \infty} \left(\frac{\alpha_k - \alpha_j}{1 - \alpha_k} \right) = 0$, for all $0 < j \leq k$;
- 3) $\lim_{n \rightarrow \infty} \beta_n = 0$;
- 4) $\sum_{n \geq 0} \beta_n = \infty$.

For an arbitrary but fixed $\mu \in C$, and for each $k \geq 0$, define $T_k: C \rightarrow C$ by $T_k x = (1 - \alpha_k)\mu + \alpha_k T x, \forall x \in C$.

Then, the double sequence $\{x_{k,n}\}_{k \geq 0, n \geq 0}$ generated from an arbitrary $x_{0,0} \in C$ by

$$\begin{aligned} x_{k,n+1} &= S y_{k,n} \\ y_{k,n} &= (1 - \beta_n)x_{k,n} + \beta_n T_k x_{k,n}, \quad k, n \geq 0, \end{aligned}$$

Converges strongly to a fixed point x_∞^* of T in C .

Proof: Consider

$$\langle T_k x - T_k y, x - y \rangle = \alpha_k \langle T x - T y, x - y \rangle \leq \alpha_k \|x - y\|^2$$



So that $\forall k \geq 0$, T_k is continuous and strongly pseudocontractive. Also, C is invariant under T_k , $\forall k$ by convexity.

Hence, T_k has a unique fixed point $x_k^* \in C$, $\forall k \geq 0$. It thus suffices to prove the following:

- i. For each fixed $\forall k \geq 0$, $x_{k,n} \rightarrow x_k^* \in C$ as $n \rightarrow \infty$;
- ii. $x_k^* \rightarrow x_\infty^* \in C$ as $k \rightarrow \infty$;
- iii. $x_\infty^* \in F(T)$.

Now, Let $b = \text{diam } C$ and $\gamma_k = (1 - \alpha_k) \in (0, 1)$, $\forall k$. Then

$$\begin{aligned} \|x_{k,n+1} - x_k^*\|^2 &= \|Sy_{k,n} - x_k^*\|^2 \leq \|Sy_{k,n} - x_k^*\|^2 \\ &= \|y_{k,n} - x_k^*\|^2 \\ &= \|(1 - \beta_n)x_{k,n} + \beta_n T_k x_{k,n} - x_k^*\|^2 \\ &= \|x_{k,n} - x_k^* - \beta_n (x_{k,n} - T_k x_{k,n})\|^2 \\ &= \|x_{k,n} - x_k^*\|^2 - 2\beta_n \langle x_{k,n} - T_k x_{k,n}, x_{k,n} - x_k^* \rangle + \beta_n^2 \|x_{k,n} - T_k x_{k,n}\|^2 \\ &= \|x_{k,n} - x_k^*\|^2 - 2\beta_n (1 - \alpha_k) \|x_{k,n} - x_k^*\|^2 + b^2 \beta_n^2 \\ &= (1 - 2\gamma_k \beta_n) \|x_{k,n} - x_k^*\|^2 + b^2 \beta_n^2 \end{aligned}$$

If we set

$$\theta_{k,n} = \|x_{k,n} - x_k^*\|, \quad \delta_{k,n} = 2\gamma_k \beta_n, \quad \sigma_{k,n} = b^2 \beta_n^2$$

Then we have

$$\theta_{k,n+1}^2 \leq (1 - \delta_{k,n}) \theta_{k,n}^2 + \sigma_{k,n},$$

So that observing that

$$\sigma_{k,n} = o(\delta_{k,n}), \quad \lim_{n \rightarrow \infty} \delta_{k,n} = 0 \quad \text{and} \quad \sum_{n \geq 0} \sigma_{k,n} = 0$$

We then have that $\theta_{k,n} \rightarrow 0$ as $n \rightarrow \infty$. So the part (i) is proved.

Now,

$$\begin{aligned} \|x_k^* - T x_k^*\| &= \|x_k^* - \alpha_k^{-1} x_k^* - \alpha_k^{-1} (1 - \alpha_k) \mu\| \\ &= \left\| \left(\frac{\alpha_k - 1}{\alpha_k} \right) x_k^* - \left(\frac{1 - \alpha_k}{\alpha_k} \right) \mu \right\| \\ &= \left(\frac{1 - \alpha_k}{\alpha_k} \right) (\|x_k^*\| + \|\mu\|) \\ &\leq 2b \left(\frac{1 - \alpha_k}{\alpha_k} \right) \end{aligned}$$

So that

$$\lim_{k \rightarrow \infty} \|x_k^* - T x_k^*\| = 0$$

And hence, $\{x_k^*\}$ is an approximate fixed point sequence for T . Also, supposing that x_∞^* is a fixed point of T , then

$$\lim_{k \rightarrow \infty} \|x_{\infty}^* - T_k x_{\infty}^*\| = \lim_{k \rightarrow \infty} (1 - \alpha_k) \|x_{\infty}^* - \mu\| \leq 2b \lim_{k \rightarrow \infty} (1 - \alpha_k) = 0$$

Now, for all $0 < j \leq k$,

$$\begin{aligned} \|x_k^* - x_j^*\|^2 &= \langle x_k^* - x_j^*, x_k^* - x_j^* \rangle \\ &= \langle T_k x_k^* - T_k x_j^*, x_k^* - x_j^* \rangle \\ &= (\alpha_j - \alpha_k) \langle \mu, x_k^* - x_j^* \rangle + \langle \alpha_k T x_k^* - \alpha_j T x_j^*, x_k^* - x_j^* \rangle \\ &= (\alpha_j - \alpha_k) \langle \mu, x_k^* - x_j^* \rangle + (\alpha_k - \alpha_j) \langle T x_j^*, x_k^* - x_j^* \rangle + \alpha_k \langle T x_k^* - T x_j^*, x_k^* - x_j^* \rangle \\ &\leq |\alpha_k - \alpha_j| \|x_k^* - x_j^*\| (\|T x_j^*\| + \|\mu\|) + \alpha_k \|x_k^* - x_j^*\|^2 \end{aligned}$$

So that

$$(1 - \alpha_k) \|x_k^* - x_j^*\|^2 \leq |\alpha_k - \alpha_j| \|x_k^* - x_j^*\| (\|T x_j^*\| + \|\mu\|)$$

And hence,

$$\lim_{k, j \rightarrow \infty} \|x_k^* - x_j^*\| \leq 2b \lim_{k, j \rightarrow \infty} \left(\frac{\alpha_k - \alpha_j}{1 - \alpha_k} \right) = 0$$

Thus, $\{x_k^*\}$ is a Cauchy sequence, and hence there exists $x_{\infty}^* \in C$ such that $x_k^* \rightarrow x_{\infty}^*$ as $k \rightarrow \infty$.

So the part (ii) is proved.

By continuity, $T x_k^* \rightarrow T x_{\infty}^*$ as $k \rightarrow \infty$. But $x_k^* - T x_k^* \rightarrow 0$ as $k \rightarrow \infty$. Hence, $x_{\infty}^* \in F(T)$.

This completes the proof.

Corollary 2.2: Let C be a bounded closed convex nonempty subset of a real Hilbert space H , and $T: C \rightarrow C$ be a continuous pseudocontractive map. Let $S: C \rightarrow C$ be nonexpansive mapping satisfying that $\|x - S y\| \leq \|S x - S y\|$ for all $x, y \in C$ and $\{\beta_n\}_{n \geq 0}, \{\alpha_k\}_{k \geq 0} \subset (0, 1)$ be real sequence satisfying the following condition :

- 1) $\lim_{k \rightarrow \infty} \alpha_k = 1$ (monotonically);
- 2) $\lim_{k, j \rightarrow \infty} \left(\frac{\alpha_k - \alpha_j}{1 - \alpha_k} \right) = 0$, for all $0 < j \leq k$;
- 3) $\lim_{n \rightarrow \infty} \beta_n = 0$;
- 4) $\sum_{n \geq 0} \beta_n = \infty$.

For an arbitrary but fixed $\mu \in C$, and for each $k \geq 0$, define $T_k: C \rightarrow C$ by $T_k x = \alpha_k T x, \forall x \in C$. Then, the double sequence $\{x_{k,n}\}_{k \geq 0, n \geq 0}$ generated from an arbitrary $x_{0,0} \in C$ by

$$\begin{aligned} x_{k,n+1} &= S y_{k,n} \\ y_{k,n} &= (1 - \beta_n) x_{k,n} + \beta_n T_k x_{k,n}, \quad k, n \geq 0, \end{aligned}$$

Converges strongly to a fixed point x_{∞}^* of T in C .

Proof: This follows from Theorem 2.1 on setting $\mu = 0 \in C$.

Corollary 2.3: In Theorem 2.1, Let T be a non expansive map. Then the same conclusion is obtained.

Proof: Observe that every non expansive map is continuous pseudocontraction.

Remark 2.5: If we put $S = I$ in Theorem 2.1 then we get the result of C. Moore (see [17])

Remark 2.4: Prototypes of the sequence $\{\alpha_k\}$ are: $\forall k \geq 0$,

$$\alpha_k = \exp\left\{-\frac{1}{k+1}\right\}, \quad \alpha_k = \log_e\left(e - \frac{1}{k+1}\right), \quad \alpha_k = 1 - \frac{1}{\log_m(k+m+1)}, \quad m > 1.$$

REFERENCES

- [1] F.E. Browder and W.V. Petryshyn, Construction of fixed points of nonlinear mappings in Hilbert spaces, *J. Math. Anal. Appl.* 20, 1967, 197-228.
- [2] R.E. Bruck, The iterative solution of the equation $y \in x + Tx$ for a monotone operator T in Hilbert spaces, *Bull. Amer. Math. Soc.* 79, 1973, 1259-1262.
- [3] C.E. Chidume, Iterative approximation of Lipschitz strictly pseudocontractive mappings, *Proc. Amer. Math. Soc.* 99, 1987, 283-288.
- [4] C.E. Chidume, Approximation of fixed points of strongly pseudocontractive mappings, *Proc. Amer. Math. Soc.* 120 (2), 1994, 545-551.
- [5] C.E. Chidume, Global iteration schemes for strongly pseudocontractive maps, *Proc. Amer. Math. Soc.* 126, 1998, 2641-2649.
- [6] C.E. Chidume, Iterative solution of nonlinear equations of strongly accretive type, *Math. Nachr.* 189, 1998, 46-60.
- [7] C.E. Chidume and C. Moore, The solution by iteration of nonlinear equations in uniformly smooth Banach spaces, *J. Math. Anal. Appl.* 215 (1), 1997, 132-146.
- [8] C.E. Chidume and C. Moore, Fixed point iteration for pseudocontractive maps, *Proc. Amer. Math. Soc.* 127 (4), 1999, 1163-1170.
- [9] C.E. Chidume and C. Moore, Steepest descent method for equilibrium points of nonlinear systems with accretive operators, *J. Math. Anal. Appl.* 245 (1), 2000, 142-160.
- [10] C.E. Chidume and M.O. Osilike, Nonlinear accretive and pseudocontractive operator equations in Banach spaces, *Nonlinear Anal. TMA* 31 (7), 1998, 779-789.
- [11] L. Deng, Iteration process for nonlinear Lipschitzian strongly secretive mappings in L_p spaces, *J. Math. Anal. Appl.* 188, 1994, 128-140.
- [12] L. Deng, On Chidume's open questions, *J. Math. Anal. Appl.* 174 (2), 1993, 441-449.
- [13] S. Ishikawa, Fixed points by a new iteration method, *Proc. Amer. Math. Soc.* 44, 1974, 147-150.

- [14] L.S. Liu, Ishikawa and Mann iteration process with errors for nonlinear strongly accretive mappings in Banach spaces, *J. Math. Anal. Appl.* 194, 1995, 114-125.
- [15] W.R. Mann, Mean value methods in iteration, *Proc. Amer. Math. Soc.* 4, 1953, 506-510.
- [16] C. Moore, The solution by iteration of nonlinear equations involving ψ -strongly accretive operators in Banach spaces, *Nonlinear Analysis* 37 (1), 1999, 125-138.
- [17] C. Moore, A Double-Sequence Iteration process for fixed points of continuous pseudocontraction, *An Int. J. Computers and Mathematics with Applications* 43, 2002, 1585-1589.
- [18] S.Gopinath, J. Gnanaraj and S.Lalithambigai, Strong convergence for Hybrid S-Iteration scheme of nonexpansive and asymptotically demicontractive mappings, *Int. J. Pure Appl. Math.*, 110(1), 2016, 153 – 163.
- [19] L. Qihou, The convergence theorems of the sequence of Ishikawa iterates for hemicontractive mappings, *J. Math. Anal. Appl.* 148, 1990, 55-62.
- [20] J. Schu, Iterative construction of fixed points of asymptotically nonexpansive mappings, *J. Math. Anal. Appl.* 158, 1991, 407-413.
- [21] H.K. Xu, Inequalities in Banach spaces with applications, *J. Nonlinear Anal. TMA* 16 (2), 1991, 1127-1138.