

Acute Toxic Stress of Copper on Blood of

Clariasbatrachus

Vikas Chandra Sharma, Dr. Raja Ram Singh

Department of Zoology, V.K.S University

ABSTRACT:

*This paper reports acute toxic stress of copper on the blood of air-breathing cat-fish *Clariasbatrachus* by using $CuSO_4 \cdot 5H_2O$ as the test chemical. The average median lethal concentration of 96hr-LC₅₀ of $CuSO_4 \cdot 5H_2O$ was calculated to be 39.50 (≈ 40.00) mg/L. The sub lethal dose i.e., 1/100th ($=0.4$ mg/L) of $CuSO_4 \cdot 5H_2O$ were selected for evaluate acute toxicity on blood of *Clariasbatrachus*.*

*Acute exposure of 0.4 mg/L of $CuSO_4 \cdot 5H_2O$ for 24hrs, 48hrs and 96hrs duration showed a highly significant decrease from 2.33 ± 0.10 , 2.12 ± 0.17 and $2.38 \pm 0.12 \times 10^6/mm^3$ respectively in comparison to control value $3.5875 \pm 0.15 \times 10^6/mm^3$ of erythrocyte in *Clariasbatrachus*. In the cases of haemoglobin, acute exposure of $CuSO_4 \cdot 5H_2O$ gave a significant decrease from 9.80 ± 0.40 , 9.67 ± 0.22 and $10.21 \pm 0.28g/100ml$ respectively when compared to its corresponding control value $10.7625 \pm 0.47g/100ml$. As far as, the variation in packed cell volume is concerned, the exposure of 0.4 mg/L of $CuSO_4 \cdot 5H_2O$ for 24hrs, 48hrs and 96hrs duration leads to a highly significant decrease from $25.33 \pm 0.56\%$, $24.17 \pm 1.00\%$ and $24.75 \pm 1.26\%$ respectively in comparison to its control value of $27.943 \pm 0.9956\%$ in *Clariasbatrachus*.*

*The observation of this study will be helpful in establishment of suitable environmental condition of *Clariasbatrachus*.*

Key words: *Copper, Clariasbatrachus, Acute exposure, Erythrocyte, Haemoglobin, Packed cell volume.*

Introduction:

Pollution is the undesirable change in the physical, chemical and biological characteristics of air, water and land that can harmfully affect the health, survival and activities of living organism. Pollution of the aquatic environment with heavy metals has become a serious health concern in recent years (World Health organization, 1993; Ansari et al, 1994). The natural ecosystems are getting extensively contaminated with heavy metals .These metals are introduced into aquatic ecosystem through various routes such as agricultural pesticides,

industrial effluents and wastes, run off, domestic garbage dumps and mining activities (Merian, 1991). Increased discharge of heavy metals into natural aquatic ecosystems can expose aquatic organisms to unnaturally high levels of these metals (Van Dyk et al, 2007). Among aquatic organisms, fish cannot escape from the hazardous effects of these pollutants, and are therefore generally considered to be the most relevant organisms for pollution monitoring in aquatic ecosystems (Van der Oost et al, 2003; Burger and Campbell, 2004). So, Fish can be used as pollution indicator.

The polluted waste water from industries and also the sewage water from domestic sources containing heavy metals find their way into the nearby water bodies. Due to their persistence and accumulative nature, aquatic environment thus, absorb heavy metals from the surrounding contaminated water which ultimately affect their health. Among these animal species, fishes are the inhabitants that cannot escape from the detrimental effects of these pollutants (Olaifa et al, 2004) and are therefore very susceptible to physical and chemical change, which may be reflected in their blood components (Wilson and Taylor, 1993). There are five potential routes for a pollutant to enter a fish. These routes are through the food, non-food particles, gills, oral consumption of water and the skin. The metal once absorbed is transported by the blood to either a storage point, such as bone or to the liver for transportation. If transported by the liver, it may be stored there, excreted in bile, or passed back into the blood for possible excretion by kidney or gills or stored in extra hepatic tissues such as fat. This is how heavy metal gets accumulated in different tissues of the fish via blood. Once heavy metals are accumulated by an aquatic organism, they can be transferred through the upper classes of the food chain and cause bio magnifications (Cumine, 1975; Mance, 1987).

Materials and Methods: In the present investigation, a fresh water air-breathing teleost, *Clarias batrachus* (Linn.) was taken as the experimental animal. *Clarias batrachus* is a carnivorous air-breathing fish belonging to the family-Claridae and order-Siluriformes.

The live fishes were collected from local fisherman of Arrah and adjacent localities during 2012-2013. The fishes were brought to the laboratory in polythene bags containing the pond water. The fishes were disinfected by washing properly with dilute KMNO₄ and then transferred to many large aquaria (90×60×45 cm). The fishes were left for acclimatization to the laboratory conditions for a fortnight. During this period water of each aquarium was changed on alternate days. The fishes were fed with pieces of goat's liver and fish food available in local market.

Fishes of each aquarium were thoroughly examined and unhealthy and injured specimens were rejected. To keep the aquaria free from fungal growth antifungal chemicals were used. The temperature of water of each aquarium was maintained at room temperature throughout the period of investigation.

Determination of 96-hrLC₅₀ Dose: The toxicity tests of the heavy metals were conducted by employing the static bioassay method as designed by Doudoroff et al, (1951) and also recommended by APHA (1989) for the toxicity test experiments. As such, two sets of experiments, each were set up to determine the LC₅₀ doses for 24hrs, 48hrs, 72hrs and 96hrs of exposure for copper.

In each set of experiment, five different concentrations of copper were taken in separate aquarium. In each concentration, ten fishes from the acclimatized fish stock were kept. The first set of experiment was conducted for 24hrs, second for 48 hrs, third for 72hrs and fourth for 96hrs respectively. The experiments were monitored round the clock and numbers of fishes died during the experiments in each concentration of each set of experiment was recorded.

The above sets of experiments were repeated with fishes formed in the present project to determine the LC₅₀ dose of copper of the experimental fish. With the help of the records of dead fishes, LC₅₀doses for 24 hrs, 48hrs, and 72hrs and 96hrs exposures for copper were determined graphically by Probit Analysis (Finney, 1971). The behaviour of the fishes on exposure to various concentrations as well as to sub lethal dose of heavy metal was observed throughout the experiment. The whole sets of experiment were repeated twice to get an average result and to minimize any error due to unavoidable reasons. Fishes were taken out for blood collection from the aquarium after the respective duration of treatment. Normal value of different blood parameters were collected before commencement of each toxicity test from the normal acclimatized fishes kept in separate aquaria and these fishes were called control groups. The values were considered as control values and were used for comparison with the experimental data's obtained from the treated fishes after conducting the toxicity test of heavy metal. After determining the sub-lethal doses of copper selected for evaluation of their toxic effect on the haematology of *Clariasbatrachus* after treatment for different duration .The median lethal concentration of 96hr -LC₅₀ of CuSo₄.5H₂O was calculated to 39.50 (\approx 40.00) mg/L. sub lethal dose i.e., 1/100th ($=$ 0.4mg/L) 96hr-LC₅₀ of CuSo₄.5H₂O were selected for acute toxicity on blood of *Clariasbatrachus*.

HAEMATOLOGICAL PARAMETERS:various haematological parameters namely total erythrocyte (RBC) count,haemoglobin (Hb)content, haematocrit value (packed cell volume or PCV) of *Clariasbatrachus* when expose to the sub lethal concentration of 0.4mg/L CuSo₄.5H₂O for 24hrs, 48hrs and 96 hrs exhibit distinct variations from the Control value of different haematological parameters. A detailed account of each parameter is given below:

TOTAL ERYTHROCYTE (RBC) COUNTS: The normal value of total erythrocyte count in the controlled *Clariasbatrachus* was $3.5875 \pm 0.15 \times 10^6 / \text{mm}^3$.The total erythrocyte count in fishes when exposed to the sub lethal concentration of copper showed following variations. The Acute exposure of 0.4 mg/L of CuSo₄.5H₂O for 24hrs, 48hrs and 96hrs duration leads to a highly significant decrease from 2.33 ± 0.10 , 2.12 ± 0.17 and $2.38 \pm 0.12 \times 10^6 / \text{mm}^3$ respectively in comparison to Control value $3.5875 \pm 0.15 \times 10^6 / \text{mm}^3$ of Erythrocyte in *Clariasbatrachus*.



HAEMOGLOBIN CONTENT: The haemoglobin content was analysed in *Clariasbatrachus* exposed to the sub lethal dose of copper for 24 hrs, 48hrs, 96hrs of exposure, a marked reduction was observed .The Acute exposure of 0.4mg/L of CuSo₄.5H₂O for 24hrs, 48hrs and 96hrs duration leads to a significant decrease from 9.80±0.40, 9.67±0.22, 10.21±0.28g/100ml respectively in comparison to Control value 10.7625±0.47g/100ml of Haemoglobin in *Clariasbatrachus*

PACKED CELL VOLUME (PCV) OR HAEMATOCRIT VALUE (HV):

The packed cell volume in the treated *Clariasbatrachus* showed a decreasing trend when exposed to different duration of copper. The Acute exposure of 0.4 mg/L of CuSo₄.5H₂O for 24hrs, 48hrs and 96hrs duration leads to a highly significant decrease from 25.33±0.56%, 24.17±1.00% and 24.75±1.26% respectively in comparison to Control value of 27.943±0.9956% in Packed cell volume in *Clariasbatrachus*.

Summary and conclusion:

The average median lethal concentration of 96 hr-LC₅₀ of CuSo₄.5H₂O calculated and its sub lethal dose selected for acute toxicity on blood of *Clariasbatrachus*.

The study showed that the effects of copper on blood of *Clariasbatrachus*. In erythrocyte it show significant decrease forAcute toxicity in comparison to their control value. In the cases of haemoglobin it show significant decrease for Acute toxicity in comparision to its control value. A significant decrease also studied in packed cell volume when compare to its control value.

Therefore changes in the haematological parameters indicate that, it may be used as indication of copper stress in *Clariasbartrachus* on exposure to elevated copper level in water bodies.Pollution and contamination of water bodies by Heavy metals like copper is harmful for environment and their related organisms and it can leads to devastating situation in environment.

REFERENCES:

Ansari AA, Singh LB and Tobschell HJ (1994): Status of anthropogenically induced metal pollution in the Kanpur Unnao industrial region of the Ganga Plain, India. Environ Geol. 38: 25-33.

APHA (1989): Standard Methods for the examinations of water and waste water. 17thed. American Public Health Association, Washington.

Burger J and CompbellRK (2004): Species differences in contaminants in fish on and adjacent to the Oak Ridge Reservation, Tennessee. Envrn. Res. 96:55.



Cumine PM (1975): Mercury levels in Goergia Otter. Mink and fresh water Fish. Bull Environ ContamToxicol. 14(2):193-196.

Doudoroff P, Anderson BG, Burdick GE, Galtsoff PS, Hart WB, Patrick R, Strong ER, Surban EW and Bon Horn WM (1951): Bioassay methods for the evaluation of acute toxicity of industrial wastes to fish. Sewage Industrial wastes. 23:1380-1397.

Finney DJ (1971): Probit analysis, III Ed. Combridge University, Press. London, New York.

Krishnamurti CR and Viswanathan P (1991): Copper in the Indian environment and its human health implication. in: Toxic metals in Indian environment . Eds: CR KrishnaMurti and PushpaViswanathan). Tata McGraw,. Hill Pub. Comp. Ltd. Pp. 188-198.

Mance G. (1987): Pollution Threat of Heavy Metals in Aquatic Environments. Elsevier Science Publishers Ltd. New York. pp. 146-151.

Merian E (1991): Metals and their Compounds in the Environment. Occurrence, AnalBiolRelev. VCH: Weinheim.

Olaifa FE, Olaifa AK, Adelaja AA, Owolavi AG (2004): Heavy metal Contamination of *Clariasgariepinus* from a lake and fish farm in Ibadan, Nigeria. Afr. J. Biomed. Res. 7:145-148.

Van Dyk JC, Pieterse M and Van Vuren JHJ (2007): Histological changes in the liver of *Oreochromismossambicus* (Cichlidae) after exposure to cadmium and Zinc. Econtoxicol Environ safe. 66:432-440.